

The critical dimension of bosonic string theory in AdS space-time

Ian Davies

Centre for Particle Theory, University of Durham, Durham DH1 3LE, UK

Email: I.J.Davies@durham.ac.uk

Paul Mansfield

Centre for Particle Theory, University of Durham, Durham DH1 3LE, UK

Email: P.R.W.Mansfield@durham.ac.uk

ABSTRACT: The Polyakov bosonic string is quantised in Euclidean Anti-de Sitter space-time using functional methods. Regularisation of the functional determinants using both heat-kernel and zeta function techniques shows that the AdS_{D+1} string reduces to the Liouville field theory, as in the flat space-time case. A Feynman graph expansion is used to evaluate (approximately) the coefficient multiplying the Liouville action; this then gives the critical dimension for this space-time as 22 (that is, 21 flat directions plus one radial direction).

KEYWORDS: Bosonic Strings, Sigma Models, AdS-CFT Correspondance.

Contents

1. Introduction	1
2. Quantisation of the bosonic string in AdS	3
2.1 Change of variables	3
2.2 Regularisation of the determinant (flat worldsheet)	4
2.3 The Weyl anomaly	6
2.4 Integrating out the f -field	10
3. Discussion	15
4. Conclusions	16
5. Appendix A: Evaluation of the heat-kernel for Ω	16
6. Appendix B: Evaluation of $\text{Det}(\Gamma^{-1})$	19

1. Introduction

It has long been suspected that there should exist a string-theoretic description of gauge theory. This idea has its origins back in the work of K. Wilson [1], where he showed that confinement in QCD could be understood within lattice gauge theory as the binding together of quarks on the lattice by colour-electric flux tubes. Unfortunately, this does not provide us with a microscopic description of the confinement mechanism in QCD, since this result is valid only in the strong coupling expansion and has no continuum limit. The large N expansion of QCD discovered by 't Hooft [2] showed that the large- N limit of gauge theories has a natural description in terms of topologically planar Feynman diagrams; this is consistent with a string-theoretic approach in terms of worldsheets. Building on these ideas, Polyakov [3] made the conjecture that there ought to be a continuum string theory description of certain gauge invariant objects in QCD, namely the Wilson loops. More recently, the AdS/CFT correspondence [4] [5] [6] has provided us with a string-theoretic description of a 4-dimensional conformal field theory with $\mathcal{N} = 4$ supersymmetry. This conjecture has been checked and confirmed in numerous papers; however, we do not yet have a concrete worldsheet realization of the correspondence. The search for such a string-theoretic formulation of QCD forms the basis for the present work.

In the long term, it is to be hoped that such an understanding will lead towards a microscopic understanding of strongly coupled QCD. In the meantime, we restrict ourselves to a more modest aim - to identify and develop the calculational techniques that may help derive the *AdS/CFT* correspondence (or some similar duality) from “first principles”.

A promising place to begin looking for a worldsheet understanding of the correspondence is the work of Polyakov [3] [7] [8]. The basic idea is to reformulate pure gauge theory entirely in terms of objects called *loop functionals*, an example of which is the Wilson loop. All equations of motion, etc., are to be rewritten in terms of functionals and operations which are defined on *loop space* - that is, the space of arbitrary continuous closed loops. The resulting equations of motion (known as the *loop equations* [9]) are then conjectured to be soluble in analogy with the Feynman integrals over trajectories in standard field theory. Instead of a sum over paths whose end points are fixed, we sum over *surfaces* whose boundaries trace out the contour on which the Wilson loop is defined. Such a sum over surfaces is precisely what we encounter in string theory. The aim, then, is to choose the string theory background in such a way that we reproduce the gauge theory equations of motion, via the loop equation. This approach to gauge fields – strings duality is much more general than *AdS/CFT* - the specific string background which we choose to use is determined by the properties of whatever gauge theory we have decided to study. In fact, Anti-de Sitter space-time is a simple example of a background which possesses *zigzag symmetry* (a property of the Wilson loop) at certain points. The claim, then, is that we should place the ends of an open string at these zigzag-symmetric points in AdS_{D+1} ; the ends of the string then trace out the contour of a Wilson loop in a D -dimensional gauge theory. In this way, we arrive at a dual description of the D -dimensional Wilson loop in terms of a $(D + 1)$ -dimensional string theory in *AdS* space-time. This construction does not use D-branes at any stage, and as such may be a way of developing an understanding of gauge fields – strings dualities for non-supersymmetric gauge theories.

It has been shown [8] that the classical minimal area in Euclidean *AdS* space approximately satisfies the loop equations for non-supersymmetric $SU(N)$ gauge theory in the large- N limit. With this in mind, it is of interest to study the quantum theory of string worldsheets in *AdS*. By using a sensible set of coordinates, we can write the Euclidean version of the AdS_{D+1} metric (which is the Lobachevsky space on the upper-half plane) in a form which is conformal to $\mathbf{R}^{D+1}$:

$$ds^2 = \frac{R^2}{y^2} \left(dy^2 + \sum_{i=1}^D dX_i^2 \right) \quad (1.1)$$

where R is the radius of the *AdS* space-time. In this paper we attempt to quantise the Polyakov bosonic string in this background. We begin by re-writing the string

partition function in a new set of $(D + 1)$ variables (section 2.1) and consider the quantisation of D of them using zeta function regularisation techniques (section 2.2). The string action is quadratic in these D fields, which correspond to the X_i 's in the metric (1.1). We initially work in the flat gauge on the worldsheet, $g_{ab} = \delta_{ab}$, and then in section 2.3 we calculate the extra terms introduced by working in the conformal gauge $g_{ab} = e^{\phi(\xi)}\delta_{ab}$ (the presence of the “Liouville mode” $\phi(\xi)$ leads to a conformal anomaly, as in the flat space-time case). Having done this, we analyse the symmetries of the resultant effective action; this then allows us to construct a Feynman integral expansion to integrate out (approximately) the remaining space-time field (section 2.4). This computation yields an approximate value for the critical dimension for bosonic strings in AdS of 22. A discussion of this result is included in section 3.

2. Quantisation of the bosonic string in AdS

2.1 Change of variables

We begin by considering the usual expression for the partition function of a bosonic string in a $(D + 1)$ -dimensional AdS space-time with Euclidean signature. This is a non-linear sigma model, with target space metric (1.1):

$$Z = \int \mathcal{D}g \mathcal{D}X \mathcal{D}y e^{-S} \quad (2.1)$$

where

$$S = \frac{R^2}{4\pi} \int d^2\xi \sqrt{g} g^{ab} \frac{\partial_a X^i \partial_b X^i}{y^2} + \frac{R^2}{4\pi} \int d^2\xi \sqrt{g} g^{ab} \frac{\partial_a y \partial_b y}{y^2} \quad (2.2)$$

Here, $i = 1, \dots, D$ and we have set $\alpha' = 1$. We do not consider any other background fields (such as the antisymmetric tensor field) here; we are only concerned with the sector of the string theory that has already been shown to satisfy the loop equations classically [7]. Functional quantisation of string theory relies on setting up all the integrals in a manifestly reparametrisation invariant way. Weyl invariance, which is a symmetry of the classical theory, is then broken quantum-mechanically through regularisation of the functional integrals. We therefore need to check by hand the conditions under which Weyl invariance is restored at the quantum level; as is explained below, this is what leads to the existence of a critical dimension in this formalism.

In order to use functional integrals to quantise the theory, we need a proper definition of the measure. Following [10], we construct the volume element in analogy with the construction of such elements in finite-dimensional integrals. This requires a definition of an inner product of worldsheet fields, and this definition must by construction respect reparametrisation invariance. The target space metric (1.1) thus dictates that the inner product on variations of the X -fields is

$$(\delta X_1^i, \delta X_2^i) \equiv \int d^2\xi \frac{\sqrt{g}}{y^2} \delta X_1^i \delta X_2^i \quad (2.3)$$

We see that the functional integration over the X -fields is a simple Gaussian, and gives

$$\text{Det}^{-\frac{D}{2}} \left[-\frac{1}{\mu^2} \frac{y^2}{\sqrt{g}} \partial_a \left(\frac{1}{y^2} \sqrt{g} g^{ab} \partial_b \right) \right] \equiv \text{Det}^{-\frac{D}{2}} \left(\frac{\Omega}{\mu^2} \right) \quad (2.4)$$

In order to proceed with the quantisation we need to regularise this infinite dimensional functional determinant, which should be thought of as an infinite product of the eigenvalues of Ω/μ^2 . Hence, we have introduced the mass scale μ^2 in order to make the determinant dimensionless. However, it is convenient to rewrite the partition function in a new set of variables,

$$W^i(\xi) = \frac{X^i(\xi)}{y(\xi)}, \quad y(\xi) = e^{\chi(\xi)} \quad (2.5)$$

and then define a new field $f(\xi)$ to be

$$f(\xi) \equiv g^{ab} \partial_a \chi \partial_b \chi + \Delta \chi \quad (2.6)$$

where Δ is the usual covariant Laplacian operator associated with the worldsheet metric g_{ab} . This puts the partition function (before integrating out any of the fields) into the form

$$Z = \int \mathcal{D}g \mathcal{D}W \mathcal{D}f \left[\text{Det}^{-1} \left(\frac{\Delta + f}{\mu^2} \right) \right] e^{-S'} \quad (2.7)$$

with

$$S' = \frac{R^2}{4\pi} \int d^2\xi \sqrt{g} (f + W \cdot (\Delta + f) \cdot W) \quad (2.8)$$

The functional determinant appearing here is the Jacobian associated with this change of variables. As described above, we must define the measure associated with the W -fields in a reparametrisation invariant way; this forces us to use the inner product

$$(\delta W_1^i, \delta W_2^i) \equiv \int d^2\xi \sqrt{g} \delta W_1^i \delta W_2^i,$$

and so the functional integration over the W -field is now itself a straightforward Gaussian. This integral gives another factor of the Jacobian determinant given above; W is a D -dimensional vector of scalar fields, and so the result of carrying out the W -integration is just

$$Z = \int \mathcal{D}g \mathcal{D}f \left[\text{Det}^{-\frac{D+2}{2}} \left(\frac{\Delta + f}{\mu^2} \right) \right] \exp \left(-\frac{R^2}{4\pi} \int d^2\xi \sqrt{g} f \right) \quad (2.9)$$

2.2 Regularisation of the determinant (flat worldsheet)

Our goal now is to regulate the infinite-dimensional determinant arising in the expression (2.9). To do this, we will begin by assuming that the string worldsheet is flat in order to extract the χ -dependence of the determinant. In this case,

$\Delta + f \rightarrow -\partial_a^2 + (\partial_a \chi)^2 - \partial_a^2 \chi$ and we can write down a *heat-kernel equation* for this operator:

$$(-\partial_a^2 + f)K(\xi, \xi'; t) = -\mu^2 \frac{\partial}{\partial t} K(\xi, \xi'; t) \quad (2.10)$$

with the initial condition

$$\lim_{t \rightarrow 0} K(\xi, \xi'; t) = \delta^2(\xi - \xi')$$

We can write the logarithm of the determinant we are interested in by using the zeta function, defined by:

$$\zeta(s) = \sum_{\lambda} \lambda^{-s}$$

where λ are the eigenvalues of $-\partial_a^2 + f$, and so

$$\ln \text{Det} \left(\frac{-\partial_a^2 + f}{\mu^2} \right) = -\zeta'(0)$$

We can represent the zeta function as an integral over the trace of $K(\xi, \xi'; t)$:

$$\zeta(s) = \frac{1}{\Gamma(s)} \int_0^\infty dt \left(t^{s-1} \text{Tr} K \right)$$

Hence, the regularisation procedure relies on knowing the heat-kernel $K(\xi, \xi'; t)$ for Ω . In order to do this, we have to make an approximation. To begin with, we can work to the leading terms in a derivative expansion of χ , so that $(\partial_a \chi)$ is constant. In this case, f itself is a positive-definite constant and as such will act as a mass term in the heat equation (2.10). We can now write down $K(\xi, \xi'; t)$ explicitly:

$$K(\xi, \xi'; t) = \frac{\mu^2}{4\pi t} e^{-\frac{\mu^2 |\xi - \xi'|^2}{4t} - \frac{ft}{\mu^2}} \quad (2.11)$$

Substituting this heat-kernel into the expression for the zeta function then gives

$$\zeta(s) = \int d^2 \xi \left[\mu^2 \left(\frac{f}{\mu^2} \right)^{1-s} \frac{1}{4\pi(s-1)} \right],$$

which then implies that

$$\ln \text{Det} \left(\frac{-\partial_a^2 + f}{\mu^2} \right) = \frac{1}{4\pi} \int d^2 \xi \left(f - f \ln \left(\frac{f}{\mu^2} \right) \right) \quad (2.12)$$

We can now substitute this determinant back into our expression for the partition function (2.9). At present, our result appears to depend explicitly on the arbitrary

mass scale μ^2 ; however, this dependence can be absorbed by a redefinition of the radius of our AdS space-time, R ¹:

$$R^2 = \frac{(D+2)}{2} \ln \left(\frac{\Lambda^2}{\mu^2} \right)$$

(Λ is a renormalisation scale, in the same sense as Λ_{QCD}). So, we arrive at the following expression for the partition function, having integrated out the W -fields within the approximation that f is constant and using a flat worldsheet metric:

$$Z = \int \mathcal{D}f \exp \left(-\frac{(D+2)}{8\pi} \int d^2\xi \left(f - f \ln \left(\frac{f}{\Lambda^2} \right) \right) \right) \quad (2.13)$$

2.3 The Weyl anomaly

An important aspect of string physics is the Weyl anomaly - in fact, in this functional formalism it is the organising principle. Reparametrisation invariance is present from the outset, but it is the restoration of Weyl invariance by hand at the quantum level that determines the consistency conditions, such as the existence of a critical dimension. For example, in flat space-time it is only in 26 dimensions that the Faddeev-Popov determinant arising from fixing the conformal gauge $g_{ab} = e^\phi \delta_{ab}$ is exactly cancelled by the functional determinant obtained by quantising the space-time fields X^μ [10]. Away from this critical dimension we are left with the *Liouville field theory*, which in the conformal gauge is

$$\int \mathcal{D}\phi \exp \left(- \int d^2\xi [(\partial_a \phi)^2 + \lambda e^\phi] \right) \quad (2.14)$$

(λe^ϕ is a finite contribution left over from the addition of a counterterm required for renormalizability). Since the functional measure $\mathcal{D}\phi$ depends on ϕ itself in a highly complicated way, this integral is intractable. The problem of solving Liouville field theory is thus central to the understanding of string theories away from their critical

¹It is perhaps worth mentioning that the above renormalisation property may have implications for AdS/CFT . If we naively apply the relation $R^2 = g_{YM}^2 N$ to the above redefinition of R , we arrive at an expression for the dual gauge theory coupling constant as a function of the scale Λ . In other words, we get a gauge theory β -function

$$\beta \equiv \Lambda \frac{dg_{YM}(\Lambda)}{d\Lambda} = \frac{(D+2)}{\sqrt{N}}$$

It is not clear what this quantity really means. Obviously we haven't even quantised all the space-time fields at this stage (and those that we have, have been calculated only approximately), so the fact that $\beta \neq 0$ should not worry us unduly. Also, the construction of Polyakov [3] is based on just a pure Yang-Mills theory, not a supersymmetric one such as $\mathcal{N} = 4$ super-Yang Mills. But it is intriguing that one side of the AdS/CFT relationship $R^2 = g_{YM}^2 N$ appears to be determined by worldsheet renormalizability in this calculation.

dimensions, since if we were able to correctly integrate out the Liouville mode, the resulting partition function would be Weyl invariant in all space-time dimensions.

Do we find the same kind of structure in *AdS* space-time? We need to determine the explicit ϕ -dependence of the determinant

$$\text{Det}(\Delta + f)$$

in order to begin to answer this question. There are several different ways of doing this; here we will use the heat-kernel expansion method used in [10]. Clearly, the determinant given in (2.4) is equivalent to $\text{Det}(\Delta + f)$, so we will compute the variation of $\ln \text{Det} \Omega$ with respect to ϕ . We can then use the result as a consistency check on the result obtained in the previous section (we should find agreement in the f -coefficient between this calculation and the result of replacing $\mu^2 \rightarrow \mu^2 e^\phi$ in (2.12) and varying it with respect to ϕ , since both μ and ϕ always appear in the combination $\mu^2 e^\phi$). In the conformal gauge, then, we are interested in

$$\text{Det} [-e^{-\phi} \partial_a^2 + 2e^{-\phi} (\partial_a \chi) \partial_a]$$

(we have dropped the μ -dependence, since we have already calculated it) and we must now use the inner product (2.3) for worldsheet scalar fields. In analogy with the identity

$$\delta \ln \det M = \int_0^\infty dt \text{tr} (\delta M e^{-tM})$$

which holds for any finite-dimensional matrix M with positive-definite eigenvalues, we can define the regularised determinant of Ω by

$$\delta_\phi \ln \text{Det} \Omega = \int_\epsilon^\infty dt \text{Tr} (\delta_\phi \Omega e^{-t\Omega}) \quad (2.15)$$

where ϵ is a short-time cutoff (we can do this, since Ω is hermitian with respect to the inner product (2.3)). Performing the t -integral, and noting that $\delta_\phi \Omega = -\delta\phi\Omega$, we obtain

$$\delta_\phi \ln \text{Det} \Omega = -\text{Tr} (\delta\phi(\xi) e^{-\epsilon\Omega}) \quad (2.16)$$

Again we make use of a heat-kernel equation, this time for Ω . We can represent $e^{-t\Omega}$ as

$$e^{-t\Omega} f(\xi) = \int d^2 \xi' e^{\phi(\xi') - 2\chi(\xi')} K(\xi, \xi'; t) f(\xi')$$

and hence

$$\frac{\partial K(\xi, \xi'; t)}{\partial t} = -\Omega K(\xi, \xi'; t)$$

with the initial condition

$$\lim_{t \rightarrow 0} K = e^{2\chi(\xi') - \phi(\xi')} \delta^2(\xi - \xi')$$

The fact that we have introduced a short-time cutoff means that we are only interested in *local* worldsheet effects. Therefore, we can suppose that the worldsheet metric we are considering is a perturbation about a flat metric, and we should be able to evaluate the heat-kernel for Ω by expanding around the heat-kernel for the flat-space Laplacian Δ . The “unperturbed” heat-kernel satisfying the heat equation for Δ is just

$$K_0 = \frac{e^{2\chi(\xi') - \phi(\xi')}}{4\pi t} e^{-\frac{|\xi - \xi'|^2}{4t}} \quad (2.17)$$

We know that

$$\int d^2\xi' e^{\phi(\xi') - 2\chi(\xi')} K_0(\xi, \xi'; t - t') K_0(\xi', \xi; t') = K_0(\xi, \xi; t) = \frac{e^{2\chi(\xi) - \phi(\xi)}}{4\pi t}$$

from looking at the expansion of the heat-kernel in eigenfunctions of its operator. Remember that the Weyl scaling behaviour of Ω is given in terms of the *trace* of its heat-kernel, so we can use the following *heat-kernel expansion* to evaluate $K(\xi, \xi; \epsilon)$:

$$\begin{aligned} K^t(\xi, \xi) &= \frac{1}{4\pi t} + \int_0^t dt' \int d^2\xi' e^{2\chi - \phi} K_{\xi\xi'}^{t-t'} V(\xi') K_{\xi'\xi}^{t'} \\ &+ \int_0^t dt' \int_0^{t'} dt'' \int d^2\xi' e^{2\chi - \phi} \int d^2\xi'' e^{2\chi - \phi} K_{\xi\xi'}^{t-t'} V(\xi') K_{\xi'\xi''}^{t'-t''} V(\xi'') K_{\xi''\xi}^{t''} \\ &+ \dots \end{aligned} \quad (2.18)$$

where we have expanded the operator Ω around the flat-space Laplacian

$$\Omega = \Delta + V(\xi)$$

and used the heat-kernel (2.17). We need to calculate all the terms in (2.18) which are of $O(1)$ in the time variable t . Terms of order t will not contribute when we send the cutoff ϵ to zero. The heat equation for K_0 implies that the heat-kernel for the operator $-e^{-\phi(\bar{\xi})} \partial_a^2$ where $e^{-\phi(\bar{\xi})}$ is a constant must be

$$\bar{K}_0(\xi, \xi'; t) = \frac{e^{2\chi(\xi') - \phi(\xi')}}{4\pi t} e^{\phi(\bar{\xi})} e^{-\frac{|\xi - \xi'|^2}{4t}} e^{\phi(\bar{\xi})} \quad (2.19)$$

and so we can write Ω as

$$\Omega = e^{-\phi(\bar{\xi})} \Delta - V(\xi)$$

where

$$\begin{aligned} V(\xi') &= \left((\xi' - \bar{\xi})^a \partial_a \phi - \frac{1}{2} (\xi' - \bar{\xi})^a (\xi' - \bar{\xi})^b (\partial_a \phi \partial_b \phi - \partial_a \partial_b \phi) \right) e^{-\phi(\bar{\xi})} \Delta' \\ &- 2 \left(1 - (\xi' - \bar{\xi})^a \partial_a \phi + \frac{1}{2} (\xi' - \bar{\xi})^a (\xi' - \bar{\xi})^b (\partial_a \phi \partial_b \phi - \partial_a \partial_b \phi) \right) \\ &\times \left(\partial_a \chi + (\xi' - \bar{\xi})^b \partial_b \partial_a \chi \right) e^{-\phi(\bar{\xi})} \partial'_a \end{aligned}$$

We are now faced with the task of substituting this expression for V into the heat-kernel expansion (2.18) and evaluating all the relevant terms, using the heat-kernel (2.19). This calculation, although now conceptually straightforward, is rather lengthy and technical. Essentially, it involves calculating a series of 2-dimensional Gaussian integrals; this is most easily done by constructing a “generating function” and taking derivatives in order to generate the required terms. For the reader’s convenience the details are given in Appendix A; here, we simply quote the result for the trace of the heat-kernel of Ω :

$$K(\xi, \xi; \epsilon) = e^{2\chi(\xi) - \phi(\xi)} \left[\frac{e^\phi}{4\pi\epsilon} - \frac{1}{24\pi} \partial_a^2 \phi + \frac{1}{4\pi} (\partial_a^2 \chi - (\partial_a \chi)^2) + O(\epsilon) \right]$$

We can now use this result to evaluate $\delta_\phi \ln \text{Det} \Omega$ by taking $\epsilon \rightarrow 0$:

$$\begin{aligned} \delta_\phi \ln \text{Det} \Omega &= \int d^2 \xi \delta \phi \left[\frac{1}{24\pi} \partial_a^2 \phi - \frac{1}{4\pi} (\partial_a^2 \chi - (\partial_a \chi)^2) \right] \\ &= \int d^2 \xi \delta \phi \left[\frac{1}{24\pi} \partial_a^2 \phi + \frac{1}{4\pi} f \right] \end{aligned} \quad (2.20)$$

We have removed the divergent term by adding a counterterm to the original string action. Notice that replacing $\mu^2 \rightarrow \mu^2 e^\phi$ in expression (2.12) and varying it with respect to ϕ gives

$$\delta_\phi \ln \text{Det} \left(\frac{-\partial_a^2 + f}{\mu^2 e^\phi} \right) = \int d^2 \xi \left[\frac{1}{4\pi} f \right] \delta \phi,$$

and we have agreement in the f -coefficients in this expression and (2.20), as expected. It is important to note that the result (2.20) is correct for *all* f - we have not made any approximation here. Therefore, we can be sure that this result captures all of the coupling between the Liouville mode $\phi(\xi)$ and $f(\xi)$.

Combining the result for the Weyl anomaly with the f -dependence derived in (2.13), we arrive at the following expression for the string partition function in AdS_{D+1} :

$$Z = \int \mathcal{D}\phi \int \mathcal{D}f \exp \left(-S' \right) \times \text{Det}_{FP}^{1/2} \quad (2.21)$$

where

$$S' = \frac{(D+2)}{8\pi} \int d^2 \xi \left(f \left(1 - \ln \left(\frac{f}{\Lambda^2} \right) + \phi \right) - \frac{1}{12} (\partial_a \phi)^2 + \lambda e^\phi \right) \quad (2.22)$$

The λe^ϕ term in the action is the finite contribution left over from the counterterm - the value of λ is fixed by some physical renormalisation condition. We have also introduced the Faddeev-Popov determinant which arises when we fix the conformal gauge on the worldsheet. This determinant was evaluated in [10] for the flat space-time case; since g_{ab} is a local quantity on the worldsheet, the Faddeev-Popov

determinant remains the same irrespective of the background space-time in which the string lives. Hence, we quote the result from [10] that

$$\text{Det}_{FP}^{1/2} = \exp \left(-\frac{13}{48\pi} \int d^2\xi [(\partial_a\phi)^2 + \lambda e^\phi] \right) = \exp \left(-\frac{13}{48\pi} S_L \right)$$

We can therefore combine the f -independent terms in (2.22) with the Liouville action arising from the Faddeev-Popov determinant to obtain

$$Z = \int \mathcal{D}\phi e^{\frac{D+2-26}{96\pi} S_L} \times \int \mathcal{D}f e^{-\frac{(D+2)}{8\pi} \int d^2\xi [f(1-\ln(\frac{f}{\Lambda^2})+\phi)]} \quad (2.23)$$

2.4 Integrating out the f -field

At first sight, integrating out f in equation (2.23) looks like an extremely difficult task. However, we can apply some simple symmetry arguments to give some clues as to how to proceed. Working for the moment in complex coordinates ($z = \xi_1 + i\xi_2$, $\bar{z} = \xi_1 - i\xi_2$) so that $f = \partial\chi\bar{\partial}\chi - \partial\bar{\partial}\chi$, we see that Z has the following conformal symmetry:

$$\delta\phi = \epsilon\partial\phi + \partial\epsilon \quad \delta\chi = \epsilon\partial\chi$$

with $\bar{\partial}\epsilon = 0$. Once the f -field has been integrated out of Z , the result will be invariant under the ϕ -transformation alone. Crucially, the Liouville action is the unique local theory without dimensionful couplings that has this symmetry. Since our derivation of the ϕ -dependence of Z used only a local cutoff, we know that the result of integrating out f must also be local, and therefore must be of the form

$$Z \sim \int \mathcal{D}\phi \exp \left(C \int d^2\xi [(\partial_a\phi)^2 + \lambda e^\phi] \right)$$

The question to be answered is: what is the overall coefficient C multiplying the Liouville action? This will determine the critical dimension for this system, since when $C = 0$ the partition function will no longer depend on the scale factor ϕ and the quantum theory will be Weyl invariant.

As was mentioned above, the calculation of the Weyl anomaly captured all of the coupling between f and ϕ . This allows us to try and integrate out f approximately using a Feynman integral-type approach. We need to find the ϕ -dependence of Z , which from equation (2.23) can be written as

$$Z = e^{\frac{D+2-26}{96\pi} S_L} \times \int \mathcal{D}f e^{-\frac{(D+2)}{8\pi} \int d^2\xi \phi f} \times \text{Det}^{-\frac{(D+2)}{2}} (-\partial_a^2 + f)$$

This then must be equal to

$$Z = \int \mathcal{D}f \text{Det}^{-N} (\Delta + f) = \int \mathcal{D}f \exp(-N \text{Tr} \ln(\Delta + f))$$

where $N = \frac{(D+2)}{2}$. We can expand the exponent,

$$\text{Tr} \ln(\Delta + f) = \text{Tr} \ln \Delta + \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \text{Tr} [(\Delta^{-1} f)^n]$$

so that

$$Z = \text{Det}^{-N} \Delta \int \mathcal{D}f \exp \left[-N \left(-\text{Tr}(\Delta^{-1} f) + \frac{1}{2} \text{Tr}(\Delta^{-1} f)^2 - \dots \right) \right]$$

We can now use the knowledge gained from our previous calculations to unpack this expression a little. The first term in the exponential ($\text{Tr}(\Delta^{-1} f)$) is actually the one which generates the $f - \phi$ coupling that we found above (a standard calculation confirms this, and checks that the coefficients in both cases are the same). The factor of $\text{Det}^{-N} \Delta$ (along with the Faddeev-Popov determinant, obviously) generates the Liouville action factor appearing in Z ,

$$\exp \left(\frac{D+2-26}{96\pi} S_L \right) \quad (2.24)$$

and does not enter into the f -integration. Rescaling f such that $\sqrt{N}f = \bar{f}$, we define Ψ to be the relevant f -integral:

$$\Psi = \int \mathcal{D}\bar{f} \exp \left[- \left(\frac{\sqrt{N}}{4\pi} \int d^2\xi \bar{f} \phi + \frac{1}{2} \text{Tr}(\Delta^{-1} \bar{f})^2 - \frac{1}{3\sqrt{N}} \text{Tr}(\Delta^{-1} \bar{f})^3 + \dots \right) \right] \quad (2.25)$$

(we will omit the bars from f from now on). We know by symmetry arguments that we are looking only for terms involving ϕ of the form $\phi \partial_a^2 \phi$, since only these are present in the Liouville action (cosmological constant-type terms of the form λe^ϕ may also arise through divergences, but these will not affect the result). Therefore, our strategy is this: we will arrange Ψ as an expansion in powers of N , where N is assumed to be large (i.e. of the order of $D \sim 26$). We then look for the contributions order-by-order in N which contain $\phi \partial_a^2 \phi$. If we work in momentum space, this involves looking for terms proportional to p^2 . These terms will be corrections to the coefficient multiplying the Liouville action (2.24). (Since we expect N to be of the order of D , terms in (2.25) of order $1/N, 1/N^2 \dots$ should be negligible. The sum of all the diagrams in the expansion will form the exponential of just the connected diagrams, in the usual way.)

We shall begin by isolating all the terms in (2.25) which are of $O(N)$ and have two powers of ϕ . There turns out to be only one; this is (writing out the various traces and operators in full)

$$\begin{aligned} O(N) : & \int \mathcal{D}f \exp \left(-\frac{1}{2} \int d^2\xi \int d^2\xi' \Delta^{-1}(\xi, \xi') f(\xi') \Delta^{-1}(\xi', \xi) f(\xi) \right) \\ & \times \frac{1}{2!(4\pi)^2} \int d^2x_1 \int d^2x_2 f(x_1) \phi(x_1) f(x_2) \phi(x_2) \end{aligned} \quad (2.26)$$

We will work in momentum space, and begin by deriving the propagator associated with the operator

$$\Delta^{-1}(\xi, \xi') \Delta^{-1}(\xi', \xi) \equiv \Gamma^{-1}(\xi, \xi')$$

This will allow us to develop a set of rules for representing this and higher-order integrals as Feynman diagrams. The Fourier transform of the ϕ -independent piece of Γ^{-1} is given by

$$I(p) = \frac{1}{(2\pi)^2} \int d^2q \frac{1}{(q^2 + m^2)} \frac{1}{((q - p)^2 + m^2)} \quad (2.27)$$

We have introduced a small mass m here to regulate the infra-red divergence of this integral. We find that

$$I(p) = \frac{1}{4\pi p^2 (\sqrt{\frac{1}{4} + \frac{m^2}{p^2}})} \times \ln \left(\frac{\sqrt{\frac{1}{4} + \frac{m^2}{p^2}} + \frac{1}{2}}{\sqrt{\frac{1}{4} + \frac{m^2}{p^2}} - \frac{1}{2}} \right) \quad (2.28)$$

which goes like

$$I(p) \sim \frac{\ln \left(\frac{p^2}{m^2} \right)}{p^2}$$

as $p/m \rightarrow \infty$. Our momentum-space propagator is then the inverse of $I(p)$, which we will denote by a wiggly line in our Feynman diagram representation:

$$\Gamma = \text{wiggly line} \sim \frac{p^2}{\ln \left(\frac{p^2}{m^2} \right)}$$

Actually, we can see immediately that the $O(N)$ contribution (2.26) will be zero, since the diagram is just



which clearly vanishes when we take the regulator m to zero at fixed momentum p .

We now go on to consider the $O(1)$ contributions. We find that there are four of them; one of which is simply the determinant $\text{Det}^{-\frac{1}{2}}(\Gamma^{-1})$, and the other three can be represented by the diagrams shown in Figure 1.

The solid lines in these diagrams represent the propagator Δ^{-1} . (For instance, the triangles appearing in the first and third diagrams represent the factors of $\text{Tr}(\Delta^{-1}f)^3$ appearing in (2.25)). The first of these diagrams appears to be proportional only to ϕ , and would seem to be zero anyway since it is not 1PI. However, the “double propagator” connecting the two bottom points of the triangle needs to be regulated in a reparametrisation invariant way, and this will introduce some extra




Figure 1: The $O(1)$ contributions to (2.25)

ϕ -dependence - we will return to this issue in a moment. To begin with, we will evaluate the determinant $\text{Det}(\Gamma^{-1})$. The details of this calculation are given in Appendix B. The result is found to be

$$\text{Det}(\Gamma^{-1}) = \exp \left(\frac{1}{12\pi} \int d^2\xi \phi(\xi) \partial_a^2 \phi(\xi) \right)$$

and hence the correction to the Liouville action factor in Z (equation (2.23)) from this determinant is

$$\frac{1}{24\pi} \int d^2\xi (\partial_a \phi)^2 \quad (2.29)$$

We now go on to consider the diagrams in Figure 1. As was mentioned above, the first diagram requires some extra regularisation, and this will introduce some extra ϕ -dependence. We replace the solid line in the double propagator with

$$G_\Delta(x, y) \equiv \int_\epsilon^\infty ds e^{-s\Delta} \delta(x - y) \frac{1}{\sqrt{g(y)}}$$

and then expand $G_\Delta(x, y)$ in powers of ϕ about flat space, as in the determinant calculation above. Analysis of the integral obtained by using this regulator, along with the other pieces of the diagram, reveals that there is an overall factor of $\frac{1}{\ln \epsilon}$ present; this clearly goes to zero when we send the cutoff to zero, so this diagram does not contribute.

The second diagram in Figure 1 represents the following asymptotic (large momentum) integral (up to factors of π , etc)

$$\sim \int d^2p \left(\frac{p^2}{\ln(\frac{p^2}{m^2})} \right)^2 \int d^2j d^2q \frac{1}{(p+q)^2 + m^2} \left(\frac{1}{q^2 + m^2} \right)^2 \frac{1}{(q+j)^2 + m^2} \frac{j^2}{\ln(\frac{j^2}{m^2})}$$

The j -integral is UV-divergent; we regulate it with a UV cutoff Λ^2 and subtract the divergence by expanding the j -integral in powers of q . What we are left with is

proportional to

$$\frac{p^2}{\ln\left(\frac{p^2}{m^2}\right)} \times \ln\left(\ln\left(\frac{\Lambda}{m}\right)\right)$$

and so this diagram also vanishes when we send the $m \rightarrow 0$ at fixed momentum.

We are left with the third diagram of Figure 1 to calculate. This diagram is in fact proportional to p^2 , and so gives a finite contribution which we need to calculate. The momentum integral is rather complicated, but can be evaluated using numerical techniques (a simple quadrature formula such as Simpson's rule will suffice). We also need to take into account the various combinatoric factors associated with the diagram, and the factors arising from the expansion of the exponential in (2.25). The total contribution of this diagram is then found to be approximately

$$\sim -0.003 \times \int d^2\xi (\partial_a \phi)^2 \quad (2.30)$$

Hence, at this order the overall coefficient multiplying the Liouville action in the string partition function is found to be

$$C = \left(\frac{D-24}{96\pi}\right) + \frac{1}{24\pi} - 0.003 \Rightarrow D_c \approx 21$$

which implies that the critical dimension for AdS_{D+1} space-time is approximately $D+1=22$. In fact, we see a slight departure from the integer value at this order in N (21.9 to 1 d.p.). Therefore, we might expect that there ought to be some non-zero contribution at the next order in N which will correct for this. Unfortunately, the $O(1/N)$ diagrams are hard to evaluate; however, we can show that there is some finite contribution.

Two of the $O(1/N)$ diagrams (see Figure 2) are actually divergent - they are proportional to $p^2 \ln(\frac{p^2}{m^2})$. Although it is difficult to evaluate either of these diagrams to check that they exactly cancel, it can be shown quite easily that both are of the form

$$p^2 \ln\left(\frac{p^2}{m^2}\right) \times \text{double integral over internal momenta}$$

The internal momentum integrands in each case can be plotted, and are found to be the same up to an overall factor which we expect to be accounted for by the combinatorics. A relative minus sign also arises from the expansion of the exponential; therefore, we expect that these two divergent diagrams cancel. We are left with three non-zero diagrams at this order, which will contribute towards the Liouville action coefficient. It is to be hoped that their contributions are small, and simply tune the result obtained at $O(1)$ towards the integer value of 22.



Figure 2: The divergent diagrams at $O(1/N)$

3. Discussion

Our calculations suggest that the critical dimension for bosonic string theory in AdS_{D+1} space-time is approximately 22 (that is, an AdS space comprising 21 X -fields and one y -field). This result differs from that derived in [11] [12], where it was found that the critical dimension is 25. Interestingly, this is the result that would be obtained if all the contributions in our Feynman diagram expansion cancelled each other. It seems unlikely that this is the case, since the $O(1)$ contribution is quite large and higher order terms are expected to be small. There is another reason to suspect that the critical dimension ought not to be 25. Going back to equation (2.7), we can represent the Jacobian $\text{Det}^{-1}(\Delta + f)$ as a functional integration over two worldsheet scalar fields v_1, v_2 :

$$\text{Det}^{-1}(\Delta + f) = \int \mathcal{D}(v_1, v_2) \exp \left(- \int d^2\xi \sqrt{g} \vec{v} \cdot (\Delta + f) \vec{v} \right)$$

So we can write Z as

$$Z = \int \mathcal{D}\psi \mathcal{D}f \exp \left(- \int d^2\xi \sqrt{g} \left[f + \vec{\psi} \cdot (\Delta + f) \vec{\psi} \right] \right)$$

where $\vec{\psi} = (v_1, v_2, \vec{W})$ is a $(D + 2)$ -dimensional vector of scalar fields for AdS_{D+1} . We can now perform the f -integral to obtain a functional delta-function,

$$Z = \int \mathcal{D}\psi \delta \left[1 + \vec{\psi} \cdot \vec{\psi} \right] \exp \left(- \int d^2\xi \sqrt{g} \vec{\psi} \cdot \Delta \vec{\psi} \right)$$

This is a non-linear sigma model with $O(D + 2)$ rotational symmetry. Notice that if the delta functional were not present this would be the partition function for a bosonic string in $(D + 2)$ -dimensional flat space, and we know that the critical dimension of this system is 26. Hence, the critical value of D itself would be 24, implying that the AdS space from which this came would have a critical dimension

of 25. However, the delta functional depends on the worldsheet metric! Thus, we cannot simply apply the argument that this is essentially the partition function for a string in $(D + 2)$ -dimensional flat space. We have computed (approximately) the explicit dependence of this delta functional on the Liouville mode, and shown it to be non-zero.

4. Conclusions

In this paper we have used several different techniques to quantise the Polyakov bosonic string in $(D + 1)$ -dimensional AdS space-time. We have shown that the Weyl structure of the AdS string is essentially the same as the string in flat space-time; the symmetries of the action obtained after quantising the $\tilde{W}$ -fields in these variables dictate that the partition function must reduce to the Liouville field theory in the conformal gauge once all the space-time fields have been integrated out. However, the critical dimension differs from the flat space-time case; our approximate calculation suggests a value of 22 for AdS , rather than the value of 26 known for flat space-time.

The fact that we have seen such a significant shift downwards in the critical dimension is perhaps an encouraging sign that the critical dimension for the superstring in this space-time may be more physically 'realistic' than 10. Ideally, we would hope to find a critical dimension of 5 for the AdS superstring, since this would correspond to a 4-dimensional dual theory in the sense of [7]. It will therefore be interesting to see the effect of introducing worldsheet supersymmetry into this calculation. (It is worth remembering that this supersymmetric worldsheet theory would still be expected to have a *non*-supersymmetric dual gauge theory description.)

In flat space, the vertex operator formalism is used to calculate string amplitudes. This formalism relies on our ability to calculate the flat space string partition function. Now that we have demonstrated the calculation of the AdS string partition function (at least approximately), it may be that some of the techniques used here will allow us to begin to compute some amplitudes, and to further investigate the relationship studied in [7], [8] between this system and the loop equations. With this in mind, the introduction of a worldsheet boundary (i.e., the extension of this work to the case of open strings) is essential, since we expect the ends of an open string to trace out the contour of the dual gauge theory Wilson loop.

5. Appendix A: Evaluation of the heat-kernel for Ω

We begin by writing down the two contributions of the correct order from the second term of (2.18):

$$\int_0^t dt' \int d^2\xi' K_{\xi\xi'}^{t-t'} e^{-\phi(\xi)} \left(-\frac{1}{2}(\xi' - \xi)^a(\xi' - \xi)^b (\partial_a \phi(\xi) \partial_b \phi(\xi) - \partial_a \partial_b \phi(\xi)) \right) \Delta' K_{\xi'\xi}^{t'} \quad (\text{A-1})$$

and

$$\int_0^t dt' \int d^2\xi' K_{\xi\xi'}^{t-t'} e^{-\phi(\xi)} 2 \left((\xi' - \xi)^a \partial_a \phi(\xi) \partial_a \chi(\xi) - (\xi' - \xi)^b \partial_b \phi(\xi) \partial_a \chi(\xi) \right) \partial'_a K_{\xi'\xi}^{t'} \quad (\text{A-2})$$

where we have chosen the constant $\bar{\xi} = \xi$. These Gaussian integrals can be evaluated using (2.19) and are found to equal

$$\frac{1}{24\pi} \left((\partial_a \phi(\xi))^2 - \partial_a^2 \phi(\xi) \right) \quad (\text{A-3})$$

and

$$\frac{1}{4\pi} \left(\partial_a^2 \chi(\xi) - \partial_a \phi(\xi) \partial_a \chi(\xi) \right) \quad (\text{A-4})$$

respectively.

We now turn to the third term of (2.18). There are four terms which contribute; the relevant pieces of $V(\xi')$ and $V(\xi'')$ respectively appearing in the integral are

$$\begin{aligned} (a) &: -2\partial_a \chi(\xi) e^{-\phi(\xi)} \partial'_a \times -2\partial_b \chi(\xi) e^{-\phi(\xi)} \partial'_b \\ (b) &: -2\partial_a \chi(\xi) e^{-\phi(\xi)} \partial'_a \times (\xi'' - \xi)^b \partial_b \phi(\xi) e^{-\phi(\xi)} \Delta'' \\ (c) &: (\xi' - \xi)^a \partial_a \phi(\xi) e^{-\phi(\xi)} \Delta' \times -2\partial_b \chi(\xi) e^{-\phi(\xi)} \partial'_b \\ (d) &: (\xi' - \xi)^a \partial_a \phi(\xi) e^{-\phi(\xi)} \Delta' \times (\xi'' - \xi)^b \partial_b \phi(\xi) e^{-\phi(\xi)} \Delta'' \end{aligned} \quad (\text{A-5})$$

These integrals, while being Gaussian, are in practice rather complicated to evaluate. Let us begin with the first of these terms, (a), which involves two single derivatives on the heat-kernels.

$$\begin{aligned} (a) &= \int_0^t dt' \int_0^{t'} dt'' \int d^2\xi' \int d^2\xi'' \frac{e^{\phi(\xi)}}{4\pi(t-t')} e^{-\frac{|\xi-\xi'|^2 e^{\phi(\xi)}}{4(t-t')}} \\ &\quad \times \partial'_a \left(\frac{e^{\phi(\xi)}}{4\pi(t'-t'')} e^{-\frac{|\xi'-\xi''|^2 e^{\phi(\xi)}}{4(t'-t'')}} \partial'_b \left[\frac{e^{\phi(\xi)}}{4\pi t''} e^{-\frac{|\xi''-\xi|^2 e^{\phi(\xi)}}{4(t'')}} \right] \right) \\ &= \int_0^t dt' \int_0^{t'} dt'' \int d^2\xi' \int d^2\xi'' \frac{e^{5\phi(\xi)}}{256\pi^3} \frac{(\xi' - \xi'')^a (\xi'' - \xi)^b}{(t-t')(t'-t'')^2(t'')^2} \\ &\quad \times e^{e^{\phi(\xi)} \left(-\frac{|\xi-\xi'|^2}{4(t-t')} - \frac{|\xi'-\xi''|^2}{4(t'-t'')} - \frac{|\xi''-\xi|^2}{4(t'')} \right)}, \end{aligned} \quad (\text{A-6})$$

all multiplied by the factor $4(\partial_a \chi)(\partial_b \chi) e^{-2\phi(\xi)}$. The simplest way to tackle these integrals is to write them in the form of a matrix equation and use a 'generating function' approach. Begin by making a change of coordinates

$$X^a = (\xi' - \xi'')^a \quad Y^a = (\xi'' - \xi)^a$$

so the ξ part of the integral becomes

$$\frac{(\partial_a \chi)(\partial_b \chi) e^{3\phi}}{64\pi^3 (t-t')(t'-t'')^2 t''^2} \int d^2 X \int d^2 Y (X)^a (Y)^b e^{e^{\phi} \left(-\frac{|X+Y|^2}{4(t-t')} - \frac{|X|^2}{4(t'-t'')} - \frac{|Y|^2}{4(t'')} \right)}$$

We can write the exponent now as *vector* $\times$ *matrix* $\times$ *vector*,

$$-\frac{1}{2} (X^1, X^2, Y^1, Y^2) \mathbf{A} (X^1, X^2, Y^1, Y^2)^T$$

where the matrix is

$$\mathbf{A} = \frac{e^\phi}{2} \begin{pmatrix} \frac{1}{t-t'} + \frac{1}{t'-t''} & 0 & \frac{1}{t-t'} & 0 \\ 0 & \frac{1}{t-t'} + \frac{1}{t'-t''} & 0 & \frac{1}{t-t'} \\ \frac{1}{t-t'} & 0 & \frac{1}{t-t'} + \frac{1}{t''} & 0 \\ 0 & \frac{1}{t-t'} & 0 & \frac{1}{t-t'} + \frac{1}{t''} \end{pmatrix}$$

Let us now define a “generating function” $Z(K)$:

$$Z(K) = \int d^2X \int d^2Y \exp \left(-\frac{1}{2} (X^a, Y^a) \mathbf{A} (X^a, Y^a)^T + K (X^a, Y^a)^T \right)$$

where K is a vector $(k_1^1, k_1^2, k_2^1, k_2^2)$. This generating function is readily evaluated to be

$$Z(K) = (\det \mathbf{A})^{-1/2} (2\pi)^2 \exp \left(\frac{1}{2} K \mathbf{A}^{-1} K^T \right) \quad (\text{A-7})$$

We see that we can now write our integral as

$$\frac{(\partial_a \chi)(\partial_b \chi) e^{3\phi}}{64\pi^3 (t-t')(t'-t'')^2 t''^2} (\partial_1)^a (\partial_2)^b Z(K)$$

evaluated at $K=0$. Here, $\partial_1^a \equiv \frac{\partial}{\partial k_1^a}$. Computing these derivatives and substituting in for t then gives

$$\begin{aligned} (a) &= \int_0^t dt' \int_0^{t'} dt'' \frac{-\delta^{ab} (\partial_a \chi)(\partial_b \chi)}{2\pi \sqrt{\frac{t^2}{t''^2 (t'-t'')^2 (t-t')^2}} (t-t')(t'-t'') t t''} \\ &= -\frac{(\partial_a \chi)^2}{4\pi} \end{aligned}$$

We now consider term (b) of (A-5). In terms of the generating function, this is

$$(b) = \int_0^t dt' \int_0^{t'} dt'' \frac{(\partial_a \phi)(\partial_b \chi) e^{3\phi}}{64\pi^3 (t-t')(t'-t'')^2 t''} (\partial_1)^a (\partial_2)^b \left(\frac{1}{t''} - \frac{\partial_2^2 e^\phi}{4t''^2} \right) Z(K)$$

It is implicitly understood that this expression is evaluated at $K=0$. This is found to be

$$\begin{aligned} (b) &= \int_0^t dt' \int_0^{t'} dt'' \frac{\delta^{ab} (\partial_a \phi)(\partial_b \chi) (t-2t'')}{2\pi \sqrt{\frac{t^2}{t''^2 (t'-t'')^2 (t-t')^2}} (t-t')(t'-t'') t^2 t''} \\ &= \frac{1}{12\pi} (\partial_a \phi)(\partial_a \chi) \end{aligned}$$

Next, we consider part (c) of (A-5). This is

$$\int_0^t dt' \int_0^{t'} dt'' \frac{(\partial_a \phi)(\partial_b \chi) e^{3\phi}}{64\pi^3 (t-t')(t'-t'')t''^2} (\partial_1 + \partial_2)^a (\partial_2)^b \left(\frac{1}{(t'-t'')} - \frac{\partial_1^2 e^\phi}{4(t'-t'')^2} \right) Z(K)$$

again setting $K = 0$. This gives

$$\begin{aligned} (c) &= \int_0^t dt' \int_0^{t'} dt'' \frac{2\delta^{ab}(\partial_a \phi)(\partial_b \chi)}{2\pi \sqrt{\frac{t^2}{t''^2(t'-t'')^2(t-t')^2}} t^2 (t'-t'')t''} \\ &= \frac{1}{6\pi} (\partial_a \phi)(\partial_a \chi) \end{aligned}$$

The last term in (A-5), (d), is

$$\frac{(\partial_a \phi)(\partial_b \phi) e^{3\phi}}{64\pi^3 (t-t')(t'-t'')t''} (\partial_1 + \partial_2)^a (\partial_2)^b \left(\frac{1}{t'-t''} - \frac{e^\phi \partial_1^2}{4(t'-t'')^2} \right) \left(\frac{1}{t''} - \frac{e^\phi \partial_2^2}{4t''^2} \right) Z(K),$$

integrated over t' and t'' . This gives

$$\begin{aligned} (d) &= - \int_0^t dt' \int_0^{t'} dt'' \frac{\delta^{ab}(\partial_a \phi)(\partial_b \phi) (t^2 - 2tt' + 6t't'' - 6t''^2)}{2\pi \sqrt{\frac{t^2}{t''^2(t'-t'')^2(t-t')^2}} t^3 (t-t')(t'-t'')t''} \\ &= - \frac{(\partial_a \phi)^2}{24\pi} \end{aligned}$$

And so finally, combining results (a), (b), (c) and (d) with (A-3) and (A-4) we find the heat-kernel for the operator Ω to be

$$K(\xi, \xi; \epsilon) = e^{2\chi(\xi) - \phi(\xi)} \left[\frac{e^\phi}{4\pi\epsilon} - \frac{1}{24\pi} \partial_a^2 \phi + \frac{1}{4\pi} (\partial_a^2 \chi - (\partial_a \chi)^2) + O(\epsilon) \right] \quad (\text{A-8})$$

6. Appendix B: Evaluation of $\text{Det}(\Gamma^{-1})$

To calculate this determinant, we begin by noting that

$$\delta_\phi \Delta^{-1} = -\Delta^{-1}(\delta_\phi \Delta) \Delta^{-1} = \Delta^{-1}(\delta\phi)$$

and so

$$\delta_\phi \left(\Delta^{-1}(\xi, \xi') \Delta^{-1}(\xi, \xi') \right) = 2 \left(\Delta^{-1}(\xi, \xi') \Delta^{-1}(\xi, \xi') \right) (\delta\phi(\xi')) \quad (\text{B-1})$$

which implies that

$$\delta_\phi \Gamma = -2(\delta\phi) \Gamma$$

Now, $\delta_\phi \ln \text{Det}(\Gamma^{-1}) = -\delta_\phi \ln \text{Det}(\Gamma)$, so we can use an identity analogous to (2.15) to obtain

$$\delta_\phi \ln \text{Det}(\Gamma^{-1}) = 2 \text{Tr}(\delta\phi) e^{-\epsilon \Gamma}$$

We now proceed as before, making an expansion in powers of ϕ about the ϕ -independent piece of Γ . If we work in momentum space we are led to consider the following expression:

$$\text{Tr}(\delta\phi)e^{-\epsilon\Gamma} = \frac{2}{(2\pi)^4} \int_0^\epsilon d\tau \int d^2q d^2p \delta\phi(-p)\phi(p) e^{-(\epsilon-\tau)\tilde{\Gamma}(p+q)} \frac{\partial}{\partial\tau} e^{-\tau\tilde{\Gamma}(p)} \quad (\text{B-2})$$

where we are using the asymptotic expression for the Fourier transform of Γ , so

$$\tilde{\Gamma}(p) \equiv \frac{p^2}{\ln\left(\frac{p^2}{m^2}\right)}$$

We can see that to obtain a non-zero result as we send the regulator $\epsilon \rightarrow 0$, we need a factor of $1/\epsilon^c$ (where c is some number) to be generated from the integral over q . Therefore, we need to study the large- q regime of this integral. We thus proceed by making an expansion for $p \ll q$, and examining the terms proportional to p^2 . We make a change of variables such that $Q = q^2$, and then put $x = Q/\ln Q$. Our integral is now

$$-\frac{2}{(2\pi)^4} \pi p^2 \int_0^\epsilon d\tau \int_0^\infty dx \frac{dQ}{dx} x e^{-\epsilon x} \left((\epsilon - \tau)\tilde{f}_1(Q) + (\epsilon - \tau)^2\tilde{f}_2(Q) \right),$$

integrated over p . The functions $\tilde{f}_1$ and $\tilde{f}_2$ are the Taylor coefficients arising from the expansion in powers of p . Integrating out τ then leaves us with an integral over x of the form

$$\int_0^\infty dx e^{-\epsilon x} (\epsilon^2 F_1(x) + \epsilon^3 F_2(x))$$

($F_1(x)$ and $F_2(x)$ are simply the Taylor coefficients, re-expressed in terms of x). Now, if we replace the lower limit in this integral with some number a , we know that the result should be independent of a as we send $\epsilon \rightarrow 0$. Thus, the integral can be computed numerically and is found to be approximately $\pi/6$. In fact, comparison with the corresponding expression for the operator Δ (which we can evaluate exactly, as in [10]) shows that the two numerical integrals obtained with this technique are the same in both cases, and therefore we know that the *exact* result must be

$$\delta_\phi \ln \text{Det}(\Gamma^{-1}) = \frac{1}{6\pi} \int d^2\xi \delta\phi(\xi) \partial_a^2 \phi(\xi) \quad (\text{B-3})$$

Acknowledgments

Ian Davies would like to thank James Gregory, Ken Lovis, Tony Padilla and David Page for helpful discussions. Ian Davies is supported by a PPARC research studentship.

References

- [1] K. G. Wilson, *Confinement of quarks*, *Phys. Rev.* **D10** (1974) 2445–2459.
- [2] G. 't Hooft, *A planar diagram theory for strong interactions*, *Nucl. Phys.* **B72** (1974) 461.
- [3] A. M. Polyakov, *String theory and quark confinement*, *Nucl. Phys. Proc. Suppl.* **68** (1998) 1, [[hep-th/9711002](#)].
- [4] J. Maldacena, *The large N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231–252, [[hep-th/9711200](#)].
- [5] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253–291, [[hep-th/9802150](#)].
- [6] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, *Gauge theory correlators from non-critical string theory*, *Phys. Lett.* **B428** (1998) 105, [[hep-th/9802109](#)].
- [7] A. Polyakov and V. Rychkov, *Gauge fields - strings duality and the loop equation*, *Nucl. Phys.* **B581** (2000) 116, [[hep-th/0002106](#)].
- [8] A. M. Polyakov and V. S. Rychkov, *Loop dynamics and AdS/CFT correspondence*, *Nucl. Phys.* **B594** (2001) 272–286, [[hep-th/0005173](#)].
- [9] Y. M. Makeenko and A. A. Migdal, *Exact equation for the loop average in multicolor QCD*, *Phys. Lett.* **B88** (1979) 135.
- [10] A. M. Polyakov, *Quantum geometry of bosonic strings*, *Phys. Lett.* **B103** (1981) 207.
- [11] H. J. de Vega and N. Sanchez, *A new approach to string quantization in curved space- times*, *Phys. Lett.* **B197** (1987) 320.
- [12] A. L. Larsen and N. Sanchez, *Mass spectrum of strings in Anti-de Sitter space-time*, *Phys. Rev.* **D52** (1995) 1051–1064, [[hep-th/9410132](#)].